

# NEW FORMULA ULTRA ROUNDUP® SUPER CONCENTRATE WEEDKILLER - 1L

Perfect for large areas

Degraded in soil by micro organisms

- **KILLS TOUGH WEEDS TO THE ROOTS**
- **TREATS UP TO 3000 m<sup>2</sup>**
- **MAKES UP TO 100 SPRAY LITRES**

Use weedkillers safely. Always read the label.

**ROUNDUP ULTRA, kills most garden weeds with a single application**

**This product is a home garden weedkiller for use on unwanted vegetation including trees and freshly cut tree stumps and areas not intended to bear vegetation (e.g. weeds growing through all hard surfaces including drives, patios, paths, tarmac, gravel and paving).**

## ROUNDUP ULTRA

ROUNDUP ULTRA is a non-selective, translocated weedkiller that is absorbed by leaves and freshly cut, woody surfaces and moves throughout the entire plant, above and below ground, to control annual and perennial weeds, including grasses and deep rooted perennial weeds. It also controls standing trees, newly cut tree stumps, tree saplings, Brambles, Bracken, Nettles, Japanese Knotweed and other tough and woody weeds. ROUNDUP ULTRA is degraded by micro-organisms in the soil.

## INSTRUCTIONS FOR USE

### When to use

Apply as a foliar spray to weeds and saplings at any time when the weeds are actively growing and have adequate leaf area to absorb the spray.

Freshly cut tree stumps and standing trees should be treated from November to March, to avoid the period of active sap flow in spring/early summer.

Avoid spraying when rain is imminent. Rain within 6 hours of spraying may reduce effectiveness, so repeat treatment may be necessary.


Avoid spraying in windy conditions, as spray drift to lawns and other wanted plants will damage or destroy them. Avoid all other spray contact with cultivated plants. If cultivated plants are treated by mistake, IMMEDIATELY AND THOROUGHLY wash off the weedkiller with ample water.

### Where to use


ROUNDUP ULTRA controls annual weeds and most perennial weeds and trees around the garden and on areas not intended to bear vegetation such as hard surfaces, patios, drives, tarmac, paths and gravel areas. It can be used to clean up weedy ground before planting vegetables, flowers, bushes and trees, and to renovate overgrown areas.

Direct sprays away from ponds and other surface water bodies. **DO NOT USE ON WANTED PLANTS. DO NOT USE ON AREAS OF LAWN UNLESS YOU WANT TO KILL THEM. DO NOT USE UNDER POLYTHENE OR GLASS**


## Where to examples




Destruction of lawn



Weeding near roses, trees and ornamentals with pressure sprayer



Clear weeds around walls, fences, trees and shrubs



Renovate overgrown areas



Clear weeds from drives and patios

## How to use

Although not a requirement, as good practice, we recommend wearing waterproof gloves and gardening footwear.

Apply by sprayer to lightly wet the leaves. 1 litre is approximately 2 pints. 5 litres is approximately 1 gallon. When diluted in 1 litre of water as surface of about 30 m<sup>2</sup> can be treated. Do not apply through, or store in, galvanised or unlined steel sprayers. Weeds show effects within 1-4 weeks. Dead or dying weeds can be dug in or composted without harming future plants.

## Annual and perennial weeds (Foliar applications)

ROUNDUP ULTRA to be added to: Sprayer - 1 litre of water		Area treated	When to treat
<b>Annual weeds:</b> Chickweed, Speedwell, Annual Meadow-grass	<b>10ml</b>	<b>30m<sup>2</sup></b>	When actively growing, usually February – November
<b>Perennial grasses:</b> Including Couch	<b>15ml</b>	<b>30m<sup>2</sup></b>	When actively growing, usually February – November
<b>Tough perennial deep rooted and broad-leaved weeds:</b> Bindweed, Dandelions, Docks, Ground Elder, Thistles (see below#)	<b>15ml</b>	<b>30m<sup>2</sup></b>	Dandelions, Docks: When actively growing, usually April – October Bindweed, Ground Elder, Thistles: around the time of flowering
Areas not intended to bear vegetation	<b>15ml</b>	<b>30m<sup>2</sup></b>	When actively growing, usually February – November
Lawn destruction	<b>20ml</b>	<b>30m<sup>2</sup></b>	Treat in a period of active growth

## Lawn destruction

Apply ROUNDUP ULTRA to actively growing lawn before the onset of dormancy, usually between May and September. Allow at least 7-10 days for ROUNDUP ULTRA to be translocated to the roots of treated lawn. Remove dying and dead lawn. Either re-seed with grass seed or re-turf according to seed or turf providers instructions. ROUNDUP ULTRA should not be used on areas of lawn unless the user wants to kill them.

## Brush weeds and saplings (Foliar applications)

ROUNDUP ULTRA to be added to: Sprayer - 1 litre of water		Area treated	Best time of year to treat
<b>Brush weeds :</b> Brambles, Nettles, Horsetail (see below*), Bracken	10ml	30m <sup>2</sup>	Brambles, Nettles, Scrub : June – September Horsetail, Bracken : July – August Fully grown Nettles : cut back in early summer and treat regrowth in September
Rhododendron	30ml	30m <sup>2</sup>	Cut back and treat re-growth in summer
<b>Saplings :</b> Beech, Sycamore, Oak, Hazel, Willow and Ash	10ml	30m <sup>2</sup>	Treat June-August, before leaf senescence

Some tough weeds (e.g. Horsetail, Ground Elder, Japanese Knotweed) may need a repeat treatment if regrowth appears.

### Freshly cut tree and shrub stumps

Cut the tree or shrub stem close to the ground in November to March, and thoroughly treat the cut surface of the stump as soon as possible after felling. Use a paint brush to apply the diluted ROUNDUP ULTRA on to the cut stump. Ensure the whole cut surface is saturated, particularly around the edges and bark of the stump. Ensure the brush is thoroughly cleaned before and after use.

**Tree stump dilutions: 20% solution in water** (e.g. a mix of 20 ml of ROUNDUP ULTRA and 80 ml water).

### Standing trees

In November to March, use an axe to cut angled notches that completely penetrate the bark of standing trees. Cut at approximately knee height, one notch for trees up to 10 cm in diameter and use two or three notches in trees over 10cm diameter. Using a ¾ or 1 inch standard paint brush, apply 2 ml (thorough wetting of the inside of the notch) per 10cm trunk diameter of neat ROUNDUP ULTRA to each cut. Do not treat in the period of active sap flow in the spring/early summer. Users should be aware that dead standing trees are a potential hazard and therefore should be felled before they fall over.

### Japanese Knotweed

Apply as a foliar spot application using a 2% solution (20 ml made up to 1 litre of water). For best results apply when the plant is flowering (usually August/September) but before autumn dieback. Good coverage is essential. Spray the underside as well as the upper surface of the leaves. Repeat treatment may be necessary if regrowth appears.

If Japanese Knotweed is growing amongst wanted plants apply 5-10 ml of a 20% (200ml ROUNDUP ULTRA made up to 1 litre of water) solution with water per stem using the following method: After mid August but before leaf fall, cut the stems of the plants approximately 200mm above the base of the cane and 40mm above a node. Rupture the central stem tissue with a screwdriver and apply 5-10ml of a 20% solution of ROUNDUP ULTRA into the hollow stem within 15 minutes of cutting using the 10ml disposable pipette supplied with the pack. Gently dispense the pipette avoiding splash back. Stems must be larger than 8mm diameter.

### Restriction

If not applied as a spot treatment, the maximum yearly dose must not exceed the dose indicated in the above crop tables. Do not walk or allow children and pets into treated areas until the product has dried on leaves to avoid transfer to lawns and other desired vegetation.

Replanting and sowing	
Light cover of annual weeds :	Replanting/sowing can commence the day after treatment.
Dense cover, including perennial weeds :	Leave at least 7 days for the weedkiller to move to underground roots and shoots before digging or cultivating soil.

After application, rinse the sprayer three times, pouring rinse water over bare ground. Triple rinse the empty bottle and pour rinse water over bare ground.

## FIRST AID

Eye contact: Immediately flush with plenty of water. If easy to do, remove contact lenses.  
 Skin contact: Take off contaminated clothing, wristwatch, jewellery. Wash affected skin with plenty of water. Wash clothes and clean shoes before re-use. Inhalation: Remove to fresh air.  
 Ingestion: Immediately offer water to drink. Never give anything by mouth to an unconscious person. Do NOT induce vomiting unless directed by medical personnel. If symptoms occur, get medical attention.

**KEEP CHILDREN AND PETS OUT OF TREATED AREAS UNTIL DRY**

## RISK AND SAFETY INFORMATION

**To avoid risks to human health and the environment, comply with the instructions for use**  
 Read label before use. • Keep out of reach of children. • Keep product away from food and drink  
 • If medical advice is needed, have product container or label at hand. • Keep product in original container. • Dispose of contents/container to a household waste recycling centre as hazardous waste except for empty containers which can be disposed of by recycling. • Contact your local council for details.

MAPP16923

PCS No. 04731

## ENVIRONMENTAL PROTECTION

**DO NOT CONTAMINATE WATER WITH PRODUCT OR ITS CONTAINER.  
 DO NOT EMPTY INTO DRAINS.**

Roundup Ultra is a soluble concentrate containing 360 g/l (28.47 %w/w) glyphosate acid (present as 441 g/l (34.86 %w/w) potassium salt of glyphosate)

Roundup® is a registered trade mark of Monsanto Technology LLC. ©Monsanto  
 Monsanto Roundup, Lawn & Garden, Salisbury House, Weyside Park, Catteshall Lane, Godalming, Surrey, GU7 1XE. Telephone: 0870 530 1010

Authorisation holder: Monsanto UK Limited, PO Box 663 Cambridge CB1 0LD, UK

Protect from frost

roundup-garden.com



Keep children and pets away from treated areas until spray has dried

